

Perfect divisibility and 2-divisibility

Maria Chudnovsky *

Princeton University, Princeton, NJ 08544, USA

Vaidy Sivaraman

Binghamton University, Binghamton, NY 13902, USA

April 25, 2017

Abstract

A graph G is said to be 2-divisible if for all (nonempty) induced subgraphs H of G , $V(H)$ can be partitioned into two sets A, B such that $\omega(A) < \omega(H)$ and $\omega(B) < \omega(H)$. A graph G is said to be perfectly divisible if for all induced subgraphs H of G , $V(H)$ can be partitioned into two sets A, B such that $H[A]$ is perfect and $\omega(B) < \omega(H)$. We prove that if a graph is (P_5, C_5) -free, then it is 2-divisible. We also prove that if a graph is bull-free and either odd-hole-free or P_5 -free, then it is perfectly divisible.

1 Introduction

All graphs considered in this article are finite and simple. Let G be a graph. The complement G^c of G is the graph with vertex set $V(G)$ and such that two vertices are adjacent in G^c if and only if they are non-adjacent in G . For two graphs H and G , H is an *induced subgraph* of G if $V(H) \subseteq V(G)$, and a pair of vertices $u, v \in V(H)$ is adjacent if and only if it is adjacent in G . We say that G *contains* H if G has an induced subgraph isomorphic to H . If G does not contain H , we say that G is *H -free*. For a set $X \subseteq V(G)$ we denote by $G[X]$ the induced subgraph of G with vertex set X . For an integer $k > 0$, we denote by P_k the path on k vertices, and by C_k the cycle on k vertices. A *path in a graph* is a sequence $p_1 - \dots - p_k$ (with $k \geq 1$) of distinct vertices such that p_i is adjacent to p_j if and only if $|i - j| = 1$. Sometimes we say that $p_1 - \dots - p_k$ *is a* P_k . A *hole* in a graph is an induced subgraph that is isomorphic to the cycle C_k with $k \geq 4$, and k is the *length* of the hole. A hole is *odd* if k is odd, and *even* otherwise. The vertices of a hole can be numbered c_1, \dots, c_k so that c_i is adjacent to c_j if and only if $|i - j| \in \{1, k - 1\}$; sometimes we write $C = c_1 - \dots - c_k - c_1$. An *antihole* in a graph is an induced subgraph that is isomorphic to C_k^c with $k \geq 4$, and again k is the *length* of the antihole. Similarly, an antihole is *odd* if k is odd, and *even* otherwise. The *bull* is the graph consisting of a triangle with two disjoint pendant edges. A graph is *bull-free* if no induced subgraph of it is isomorphic to the bull. The chromatic number of a graph G is denoted by $\chi(G)$ and the clique number by $\omega(G)$. A graph G is called *perfect* if for every induced subgraph H of G , $\chi(H) = \omega(H)$. For a set X of vertices, we will usually write $\chi(X)$ instead of $\chi(G[X])$, and $\omega(X)$

*Partially supported by NSF grant DMS-1550991 and by US Army Research Office grant W911NF-16-1-0404.

instead of $\omega(G[X])$. If X is a set of vertices and x is a vertex, we will write $X + x$ for $X \cup \{x\}$.

A graph G is said to be *2-divisible* if for all (nonempty) induced subgraphs H of G , $V(H)$ can be partitioned into two sets A, B such that $\omega(A) < \omega(H)$ and $\omega(B) < \omega(H)$. Hoàng and McDiarmid [5] defined the notion of 2-divisibility. They actually conjecture that a graph is 2-divisible if and only if it is odd-hole-free. A graph is said to be *perfectly divisible* if for all induced subgraphs H of G , $V(H)$ can be partitioned into two sets A, B such that $H[A]$ is perfect and $\omega(B) < \omega(H)$. Hoàng [4] introduced the notion of perfect divisibility and proved ([4]) that (banner, odd hole)-free graphs are perfectly divisible. A nice feature of proving that a graph is perfectly divisible is that we get a quadratic upper bound for the chromatic number in terms of the clique number. More precisely:

Lemma 1.1. Let G be a perfectly divisible graph. Then $\chi(G) \leq \binom{\omega(G)+1}{2}$.

Proof. Induction on $\omega(G)$. Let $\omega(G) = \omega$. Let $X \subseteq V(G)$ such that $G[X]$ is perfect and $\chi(G \setminus X) < \omega$. Since $G \setminus X$ is perfectly divisible, $\chi(G \setminus X) \leq \binom{\omega}{2}$. Since $G[X]$ is perfect, $\chi(X) \leq \omega$. Consequently, $\chi(G) \leq \chi(G \setminus X) + \chi(X) \leq \omega + \binom{\omega}{2} = \binom{\omega+1}{2}$. \square

Analogously, 2-divisibility gives an exponential χ -bounding function.

Lemma 1.2. Let G be a 2-divisible graph. Then $\chi(G) \leq 2^{\omega(G)-1}$.

Proof. Induction on $\omega(G)$. Let $\omega(G) = \omega$. Let (A, B) be a partition of $V(G)$ such that $\omega(A) < \omega$ and $\omega(B) < \omega$. Now $\chi(A) \leq 2^{\omega-2}$ and $\chi(B) \leq 2^{\omega-2}$. Consequently, $\chi(G) \leq \chi(A) + \chi(B) \leq 2^{\omega-2} + 2^{\omega-2} = 2^{\omega-1}$. \square

We end the introduction by setting up the notation that we will be using. For a vertex v of a graph G , $N(v)$ will denote the set of neighbors of v (we write $N_G(v)$ if there is a risk of confusion). The closed neighborhood of v , denoted $N[v]$, is defined to be $N(v) + v$. We define $M(v)$ (or $M_G(v)$) to be $V(G) \setminus N[v]$. Let X and Y be disjoint subsets of $V(G)$. We say X is complete to Y if every vertex in X is adjacent to every vertex in Y . We say X is anticomplete to Y if every vertex in X is non-adjacent to every vertex in Y . A set $X \subseteq V(G)$ is a *homogeneous set* if $1 < |X| < |V(G)|$ and every vertex of $V(G) \setminus X$ is either complete or anticomplete to X . If G contains a homogeneous set, we say that G *admits a homogeneous set decomposition*.

This paper is organized as follows. In section 2 we prove that if a graph contains neither a P_5 nor a C_5 , then it is 2-divisible. In Section 3 we prove that if a graph is bull-free and either odd-hole-free or P_5 -free, then it is perfectly divisible.

2 (P_5, C_5) -free graphs are 2-divisible

We start with some definitions. Let G be a graph. $X \subseteq V(G)$ is said to be *connected* if $G[X]$ is connected, and *anticonnected* if $G^c[X]$ is connected. For $X \subseteq V(G)$, a *component* of X is a maximal connected subset of X , and an *anticomponent* of X is a maximal anticonnected subset of X .

The following lemma is used several times in the sequel.

Lemma 2.1. Let G be a graph. Let $C \subseteq V(G)$ be connected, and let $v \in V(G) \setminus C$ such that v is neither complete nor anticomplete to C . Then there exist $a, b \in C$ such that $v - a - b$ is a path.

Proof. Since v is neither complete nor anticomplete to C , it follows that both the sets $N(v) \cap C$ and $M(v) \cap C$ are non-empty. Since C is connected, there exist $a \in N(v) \cap C$ and $b \in M(v) \cap C$ such that $ab \in E(G)$. But now $v - a - b$ is the desired path. This completes the proof. \square

We are ready to prove the main result of this section.

Theorem 2.1. Every (P_5, C_5) -free graph is 2-divisible.

Proof. Let G be a (P_5, C_5) -free graph. We may assume that G is connected. Let $v \in V(G)$, let $N = N(v)$, $M = M(v)$. Let C_1, \dots, C_t be the components of M .

(1) We may assume that there is i such that no vertex of N is complete to C_i .

For, otherwise, $X_1 = M + v$, $X_2 = N$ is the desired partition. This proves (1).

Let i be as in (1), we may assume that $i = 1$.

(2) There do not exist n_1, n_2 in N and m_1, m_2 in M such that n_1 is adjacent to m_1 and not to m_2 , and n_2 is adjacent to m_2 and not to m_1 , and n_1 is non-adjacent to n_2 .

For, otherwise, $G[\{n_1, n_2, m_1, m_2, v\}]$ is a P_5 or a C_5 . This proves (2).

(3) For every $i > 1$ there exists $n \in N$ complete to C_i .

For suppose that there does not exist $n \in N$ that is complete to C_2 . For $i = 1, 2$ let $n_i \in N$ have a neighbor in C_i . Since C_1, C_2 are connected, by Lemma 2.1, there exist $a_i, b_i \in C_i$ such that $n_i - a_i - b_i$ is a path. Since $b_1 - a_1 - n_1 - a_2 - b_2$ is not a P_5 , we deduce that $n_1 \neq n_2$, and therefore n_1 is complete or anticomplete to C_2 , and n_2 is complete or anticomplete to C_1 . By the choice of C_1 and the assumption, n_1 is anticomplete to C_2 , and n_2 to C_1 . By (2) n_1 is adjacent to n_2 . But now $b_2 - a_2 - n_2 - n_1 - a_1$ is a P_5 , a contradiction. This proves (3).

From the set of vertices in N that have a neighbor in C_1 , choose one that has the maximum number of neighbors in M ; call it n . (Such a vertex exists because G is connected.) Let $X_1 = N(n)$, and let $X_2 = V(G) \setminus X_1$. Clearly X_1 does not contain a clique of size $\omega(G)$. We claim that $\omega(X_2) < \omega(G)$, thus proving that (X_1, X_2) is a partition certifying 2-divisibility.

Suppose that there is a clique K of size $\omega(G)$ in X_2 . Then $n \notin K$. By (3), $K \setminus (C_2 \cup \dots \cup C_t) \neq \emptyset$.

(4) $K \not\subseteq C_1$.

For suppose that $K \subseteq C_1$. Then $K \subseteq C_1 \setminus N(n)$. Let D be the component of $C_1 \setminus N(n)$ containing K . Then some vertex $p \in N(n) \cap C_1$ has a neighbor in D . Since D contains a clique of size $\omega(G)$, p is not complete to D . Since D is connected, by Lemma 2.1, there exist $d_1, d_2 \in D$ such that $p - d_1 - d_2$ is a path. But now $d_2 - d_1 - p - n - v$ is a P_5 , a contradiction. This proves (4).

It follows from (4) that K has a vertex $k_1 \in N \setminus X_1$, and a vertex $k_2 \in M \setminus X_1$. Then k_1 is non-adjacent to n , and k_2 is non-adjacent to n . But now by (2) $N(k_1) \cap M$ strictly contains $N(n) \cap M$, and in particular k_1 has a neighbor in C_1 , contrary to the choice of n . This completes the proof. \square

An easy consequence of this is

Corollary 2.1. Let G be a (P_5, C_5) -free graph. Then $\chi(G) \leq 2^{\omega(G)-1}$.

Proof. Follows from Theorem 2.1 and Lemma 1.2 \square

3 Perfect divisibility in bull-free graphs

For an induced subgraph H of a graph G , a vertex $c \in V(G) \setminus V(H)$ that is complete to $V(H)$ is called a *center* for H . Similarly, a vertex $a \in V(G) \setminus V(H)$ that is anticomplete to $V(H)$ is called an *antcenter* for H . For a hole $C = c_1 - c_2 - c_3 - c_4 - c_5 - c_1$, an *i-clone* is a vertex adjacent to c_{i+1} and c_{i-1} , and not to c_{i+2}, c_{i-2} (in particular c_i is an *i-clone*). An *i-star* is a vertex complete to $V(C) \setminus c_i$, and non-adjacent to c_i . A *clone* is a vertex that is an *i-clone* for some i , and a *star* is a vertex that is an *i-star* for some i . We will need the following results from [2] and [3].

Theorem 3.1. (from [3]) If G is bull-free, and G has a P_4 with a center and an antcenter, then G admits a homogeneous set decomposition, or G contains C_5 .

Theorem 3.2. (from [2]) If G is bull-free and contains an odd hole or an odd antihole with a center and an antcenter, then G admits a homogeneous set decomposition.

Theorem 3.3. (from [2]) If G is bull-free, then either G admits a homogeneous set decomposition, or for every $v \in V(G)$, either $G[N(v)]$ or $G[M(v)]$ is perfect.

The next two theorems refine Theorem 3.3 in the special cases we are dealing with in this paper.

Theorem 3.4. If G is bull-free and odd-hole-free, then either G admits a homogeneous set decomposition, or for every $v \in V(G)$ the graph $G[M(v)]$ is perfect.

Proof. We may assume that G does not admit a homogeneous set decomposition. Let $v \in V(G)$ such that $G[M(v)]$ is not perfect. Since G is odd-hole-free, by the strong perfect graph theorem [1], $G[M(v)]$ contains an odd antihole of length at least 7, and therefore a three-edge-path P with a center. Now v is an antcenter for P , and so by Theorem 3.1, G admits a homogeneous set decomposition, a contradiction. This proves the theorem. \square

Theorem 3.5. If G is bull-free and P_5 -free, then either G admits a homogeneous set decomposition, or for some $v \in V(G)$, $G[M(v)]$ is perfect.

Proof. By Theorem 3.4 we may assume that G contains a C_5 , say $C = c_1 - c_2 - c_3 - c_4 - c_5 - c_1$. We may assume that G does not admit a homogeneous set decomposition.

(1) Let D be a hole of length 5, and let $v \notin V(D)$. Then v is a clone, a star, a center or an antcenter for D .

Since G has no P_5 , v cannot have exactly one neighbor in D . Suppose that v has exactly two neighbors in D . Since G is bull-free, the neighbors are non-adjacent, so v is a clone. Suppose that v has exactly two non-neighbors in D . Since G is bull-free, the non-neighbors are adjacent, and v is a clone. The cases when v has 0, 4, 5 neighbors in D result in v being an anticenter, star, and a center for D , respectively. This proves (1).

(2) Let D be a hole of length 5 in G . Then there is no anticenter for D .

Suppose that v is an anticenter for D , we may assume that $D = C$. By Theorem 3.3 there is no center for D . Since G is connected, we may assume that v has a neighbor u such that u has a neighbor in $V(D)$. Let P be a path starting at u and with $V(P) \setminus u \subseteq V(D)$ with $|V(P)|$ maximum. Since $v - u - P$ is not a P_5 , and v is not a center for P , it follows that for some i , v is adjacent to c_i and to c_{i+1} , but not to c_{i+2} . But now $G[\{c_i, c_{i+1}, c_{i+2}, u, v\}]$ is a bull, a contradiction. This proves (2).

(3) Let d_i and d'_i be i -clones non-adjacent to each other. Let v be adjacent to d_i and not to d'_i . Then v is a center for C , or v is an i -star for C , or v is an i -clone for C . Moreover, let D be the hole obtained from C by replacing c_i with d_i , and let D' be the hole obtained from C by replacing c_i with d'_i . It follows that either

- v is an i -clone for both D and D' , or
- v is a center for D , and an i -star for D' .

We may assume that $i = 1$. If v is anticomplete to $\{c_2, c_5\}$, then we get a contradiction to (1) or (2) applied to v and D' . Thus we may assume that v is adjacent to c_2 . Suppose that v is non-adjacent to c_5 . By (1) applied to D , v is adjacent to c_3 . But now $d'_1 - c_5 - d_1 - v - c_3$ is a P_5 , a contradiction. Thus v is adjacent to c_5 . By (1) applied to D' , v is either complete or anticomplete to $\{c_3, c_4\}$. Now if v is anticomplete to $\{c_3, c_4\}$, then v is an i -clone; if v is complete to $\{c_3, c_4\}$ then v is a center or an i -star for C . This proves (3).

(4) There do not exist $d_1, d'_1, d_3, d'_3, v_1, v_3$ such that

- $\{d_1, d'_1\}$ is not complete to $\{d_3, d'_3\}$, and
- for $i = 1, 3$
 - d_i and d'_i are i -clones non-adjacent to each other, and
 - v_i is adjacent to d_i and non-adjacent to d'_i , and
 - v_i is not an i -clone.

Observe that by (3), no vertex of $\{d_1, d'_1\}$ is mixed on $\{d_3, d'_3\}$ and the same with the roles of 1, 3 exchanged. It follows that $\{d_1, d'_1\}$ is anticomplete to $\{d_3, d'_3\}$, and in particular $v_1, v_3 \notin \{d_1, d'_1, d_3, d'_3\}$. By (3) applied to the hole $d'_1 - c_2 - c_3 - c_4 - c_5 - d'_1$ and d_3, d'_3 , it follows that v_3 is complete to $\{d_1, d'_1\}$. Similarly v_1 is complete to $\{d_3, d'_3\}$. In particular $v_1 \neq v_3$. But now $G[\{d'_1, v_3, d_1, v_1, d'_3\}]$ is either a bull or a P_5 , in both cases a contradiction. This proves (4).

(5) There is not both a 1-clone non-adjacent to c_1 , and a 3-clone non-adjacent to c_3 .

For suppose that such clones exist. For $i = 1, 3$ let X_i be a maximal anticonnected set of i -clones with c_i in X_i . Then $|X_i| > 1$ for $i = 1, 3$. Since X_i is anticonnected, it follows from (3) that X_1 is anticomplete to X_3 . Since $|X_1|, |X_3| > 1$, and G does not admit a homogeneous set decomposition, it follows that neither X_1 nor X_3 is a homogeneous set in G . Therefore for $i = 1, 3$ there exists $v_i \notin X_i$ with a neighbor and a non-neighbor in X_i . Then $v_i \notin X_1 \cup X_3$. Note that $X_i + v_i$ is anticonnected, and hence by the maximality of X_i , it follows that v_i is not an i -clone. By applying Lemma 2.1 in G^c with v_i and X_i for $i = 1, 3$, it follows that there exist $d_i, d'_i \in X_i$ such that d_i is non-adjacent to d'_i , v_i is adjacent to d_i , and v_i is non-adjacent to d'_i . But now we get a contradiction to (4). This proves (5).

(6) For some i , $V(G) = N[c_i] \cup N[c_{i+2}]$ (here addition is *mod* 5).

Suppose that (6) is false. Since (6) does not hold with $i = 1$, (1), (2) and symmetry imply that we may assume that there is a 1-clone c'_1 non-adjacent to c_1 . Since (6) does not hold with $i = 5$, again by (1), (2) and symmetry we may assume that there is a 2-clone c'_2 non-adjacent to c_2 . Finally, since (6) does not hold with $i = 3$, by (1), (2) and symmetry we get a 3-clone c'_3 non-adjacent to c_3 . But this is a contradiction to (5). This proves (6).

Let i be as in (6); we may assume that $i = 1$. Suppose that $G[M(c_1)]$ is not perfect. Then, by the strong perfect graph theorem [1], $G[M(c_1)]$ contains an odd hole or an odd antihole H . But now c_3 is a center for H , and c_1 is an anticenter for H , contrary to Theorem 3.2. This proves the theorem. \square

A graph G is *perfectly weight divisible* if for every non-negative integer weight function w on $V(G)$, there is a partition of $V(G)$ into two sets P, W such that $G[P]$ is perfect and the maximum weight of a clique in $G[W]$ is smaller than the maximum weight of a clique in G .

Theorem 3.6. A minimal non-perfectly weight divisible graph does not admit a homogeneous set decomposition.

Proof. Let G be such that all proper induced subgraphs of G are perfectly weight divisible. Let w be a weight function on $V(G)$. Let X be a homogeneous set in G , with common neighbors N and let $M = V(G) \setminus (X \cup N)$. Let G' be obtained from G by replacing X with a single vertex x of X with weight $w(x)$ equal to the maximum weight of a clique in $G[X]$. Let T be the maximum weight of a clique in G .

Let (P', W') be a partition of $V(G')$ corresponding to the weight w . Let (X_p, X_w) be a partition of X where $G[X_p]$ is perfect and the maximum weight of a clique in $G[X_w]$ is smaller than the maximum weight of a clique in $G[X]$. We construct a partition of $V(G)$.

Suppose first that $x \in W'$. Then let $P = P'$ and $W = W' \cup X$. Clearly this is a good partition. Now suppose that $x \in P'$. Let $P = (P' \setminus x) \cup X_p$ and let $W = W' \cup X_w$. By a theorem of [6], $G[P]$ is perfect. Suppose that W contains a clique K with weight T . Then $K \cap X_w$ is non-empty. Let K'

be a clique of maximum weight in X . Now $(K \setminus X_w) \cup K'$ is a clique in G with weight greater than T , a contradiction. This proves the theorem. \square

We can now prove our main result:

Theorem 3.7. Let G be a bull-free graph that is either odd-hole-free or P_5 -free. Then G is perfectly weight divisible, and hence perfectly divisible.

Proof. Let G be a minimal counterexample to the theorem. Then there is a non-negative integer weight function w on $V(G)$ for which there is no partition of $V(G)$ as in the definition of being perfectly weight divisible. Let U be the set of vertices of G with $w(v) > 0$, and let $G' = G[U]$. By theorems 3.4, 3.5, 3.6, G' has a vertex v such that $G'[M_{G'}(v)]$ is perfect. But now, since $w(v) > 0$, setting $P = M_{G'}(v) + v$ and $W = N_{G'}(v) \cup (V(G) \setminus U)$ we get a partition of $V(G)$ as in the definition of being perfectly weight divisible, a contradiction. This proves the theorem. \square

Corollary 3.1. Let G be a bull-free graph that is either odd-hole-free or P_5 -free. Then $\chi(G) \leq \binom{\omega(G)+1}{2}$.

Proof. Follows from Theorem 3.7 and Lemma 1.1. \square

4 Acknowledgment

This research was performed during the 2017 Barbados Graph Theory Workshop at the McGill University Bellairs Research Institute in Barbados, and the authors are grateful to the institute for its facilities and hospitality. The authors also thank Chinh T. Hoàng for telling them about these problems, and for many useful discussions.

References

- [1] M. Chudnovsky, N. Robertson, P. Seymour, R. Thomas, The strong perfect graph theorem, *Annals of Mathematics*, 164 (2006), 51-229 .
- [2] M. Chudnovsky, S. Safra, The Erdős-Hajnal Conjecture for bull-free graphs, *Journal of Combinatorial Theory, Ser. B*, 98 (2008), 1301-1310.
- [3] M. Chudnovsky, V. Sivaraman, Odd holes in bull-free graphs, submitted.
- [4] C. T. Hoàng, On the structure of (banner, odd hole)-free graphs, submitted. (Available at <https://arxiv.org/abs/1510.02324>)
- [5] C. T. Hoàng, C. McDiarmid, On the divisibility of graphs, *Discrete Math.* 242 (2002), no. 1-3, 145-156.
- [6] L. Lovász, Normal hypergraphs and the perfect graph conjecture, *Discrete Math.* 2 (1972), no. 3, 253-267.